



Experience-based and evidence-oriented
Corticobasal® Implantology

Vol. 20 No. 8

August 2026

English Edition



FULL-MOUTH IMMEDIATE REHABILITATION WITH BASAL
(CORTICAL) IMPLANTS COMBINED WITH SURGICAL
CORRECTION OF BIMAXILLARY DENTAL PROTRUSION:
A CASE REPORT

DR. SALEM BARMAWI, DR. AMINA SALEH ABDULBAKI

ISSN 1864-1199 / e-ISSN 1864-1237

OFFPRINT

Published by IF® Publishing, Germany,
continued since 2024 as:

- **CMF Implant Directions® (CMF)**
- **Journal of unwanted Results (JUR)**
- **Experience-based and evidence-oriented
Corticobasal® Implantology (EECI)**



The Foundation of Knowledge

Editorial Board

Editor-in-Chief

Prof. Dr. Stefan Ihde
prof@ihde.com

Coordinating Editor

Prof. Dr. Fadia Awadalkreem, Dubai
editor@implantfoundation.org

Editorial Board (in Alphabetic Order)

Dr. Ashish Chakranarayan, India
Dr. Anita Doshi, India
Prof. Dr. Antonina Ihde, Belarus
Prof. Dr. Vitomir S. Konstantinovic, Serbia
Prof. Dr. Aleksandar Lazarov, Bulgaria
Dr. Juri Mitrushchenkov, Russia
Dr. Mahendra Perumal, India
Prof. Dr. Olga Sipić, Serbia
Prof. Dr. Jan Vares, Ukraine

Aspirants to the Editorial Board (in Alphabetic Order)

Dr. Salem Barmawi, Libya
Dr. Pablo Diaz, Ecuador
Dr. Marcos Daniel Gonzales, Colombia
Dr. Georg Huber, Germany
Dr. Mehul D Jani, India
Dr. Valeri Lysenko, Ukraine
Dr. Kiran Patel, India
Dr. Faizur Rahman, India
Dr. Egammai Sethuraman, India
Dr. Nikolai Spiridonov, Russia

Evidence Reports and Critical Appraisals

IF® Research & Evidence Dept.

Annual Subscription

Euro 4.600

Copyright

Copyright © 2006 - 2026 by
International Implant Foundation IF®
DE- 80802 Munich / Germany
www.implantfoundation.org

Contact

publishing@implantfoundation.org

EECI

ISSN 1864-1199
e-ISSN 1864-1237

Disclaimer

Hazards

Great care has been taken to maintain the accuracy of the information contained in this publication. However, the publisher and / or the distributor and / or the editors and / or the authors cannot be held responsible for errors or any consequences arising from the use of the information contained in this publication. The statements or opinions contained in editorials and articles in this publication are solely those of the authors thereof and not of the publisher, and / or the distributor, and / or the IF®.

The products, procedures and therapies described in this work are hazardous and are therefore only to be applied by certified and trained medical professionals in environment specially designed for such procedures. No suggested test or procedure should be carried out unless, in the user's professional judgment, its risk is justified. Whoever applies products, procedures and therapies shown or described in this publication will do this at their own risk. Because of rapid advances in the medical science, the IF® recommends that independent verification of diagnosis, therapies, drugs, dosages and operation methods should be made before any action is taken.

Although all advertising material which may be inserted into the work is expected to conform to ethical (medical) standards, inclusion in this publication does not constitute a guarantee or endorsement by the publisher regarding quality or value of such product or of the claims made of it by its manufacturer.

Legal restrictions

This work was produced by IF® Publishing, Munich, Germany. All rights reserved by IF® Publishing. This publication including all parts thereof, is legally protected by copyright. Any use, exploitation or commercialization outside the narrow limits set forth by copyright legislation and the restrictions on use laid out below, without the publisher's consent, is illegal and liable to prosecution. This applies in particular to photostat reproduction, copying, scanning or duplication of any kind, translation, preparation of microfilms, electronic data processing, and storage such as making this publication available on Intranet or Internet.

Some of the products, names, instruments, treatments, logos, designs, etc. referred to in this publication are also protected by patents and trademarks or by other intellectual property protection laws (eg. "IF®", "IIF" and the IF® Logo) are registered trademarks even though specific reference to this fact is not always made in the text.

Therefore, the appearance of a name, instrument, etc. without designation as proprietary is not to be construed as a representation by publisher that it is in the public domain.

Institutions' subscriptions allow to reproduce tables of content or prepare lists of articles including abstracts for internal circulation within the institutions concerned. Permission of the publisher is required for all other derivative works, including compilations and translations. Permission of the publisher is required to store or use electronically any material contained in this journal, including any article or part of an article. For inquiries contact the publisher at the address indicated.



FULL-MOUTH IMMEDIATE REHABILITATION WITH BASAL (CORTICAL) IMPLANTS COMBINED WITH SURGICAL CORRECTION OF BIMAXILLARY DENTAL PROTRUSION: A CASE REPORT

Authors

Dr. Salem Barmawi ^{1, 2}

Dr. Amina Saleh Abdulbaki ²

¹ Master of Science in Corticobasal® Implantology

² Misk Dental Center, Tripoli, Libya

Corresponding Author:

Dr. Salem Barmawi

drsalem.sb@gmail.com

How to Cite This Article

Barmawi S, Abdulbaki A S. Full-Mouth Immediate Rehabilitation with Basal (Cortical) Implants Combined with Surgical Correction of Bimaxillary Dental Protrusion: A Case Report, Experience-based and evidence-oriented Corticobasal® Implantology (EECI), Vol. 20, No. 8:3-20, International Implant Foundation Publishing, 2026

Abstract

Background: Bimaxillary dental protrusion combined with posterior tooth loss creates a complex esthetic, functional, and psychological burden for affected patients. Conventional correction typically requires prolonged orthodontic treatment, often combined with premolar extractions or orthognathic surgery, followed by a separate implant-prosthetic phase to replace missing posterior teeth. Basal (cortical) implant systems, which anchor into resorption-resistant cortical bone and permit immediate functional loading, offer an alternative that allows simultaneous correction of alveolar protrusion and full-arch rehabilitation within a single, compressed treatment timeline.

Case Presentation: We report the case of a 57-year-old female patient who presented with severe bimaxillary anterior dental protrusion accompanied by multiple missing posterior teeth in both the maxilla and the mandible. The pronounced dentoalveolar protrusion had resulted in advanced gingival recession, with several teeth displaced beyond the confines of the alveolar housing due to excessive labial inclination and protrusion. The extensive posterior tooth loss,

together with the advanced dentoalveolar changes, had led to a loss of the vertical dimension of occlusion (VDO), resulting in an incorrect maxillo-mandibular relationship and significant occlusal dysfunction. The patient reported severe psychological distress, including repeated bullying and mockery from family members and people around her because of her dental and facial appearance, and requested a rapid, comprehensive solution.

Following clinical and radiographic (OPG) assessment, a combined surgical-prosthetic protocol was carried out in a single treatment episode. All remaining compromised and malpositioned teeth were extracted, and approximately 7 mm of protrusive alveolar bone was resected from the maxilla to correct the dentoalveolar profile and expose a level cortical bed. Ten immediate Corticobasal® implants (Manufacturer: Dr. Ihde Dental AG, Dorfplatz 11, 8737 Gommiswald, Switzerland) were then placed in the maxilla. The identical protocol extraction, alveolar bone resection, and implant placement was subsequently carried out in the mandible, where eight additional basal implants were placed, for a total of

eighteen basal implants across both arches. Treatment continued with metal framework try-in, occlusal verification, and delivery of the final fixed prosthesis.

Conclusion: Combining surgical reduction of the protrusive alveolar segment with immediate basal implant placement allows simultaneous correction of bimaxillary protrusion and full-arch rehabilitation of missing posterior teeth in a markedly shortened treatment timeline compared to conventional orthodontic-surgical implant sequencing. In a patient whose primary complaint was esthetic and psychosocial, the rapid transformation of facial profile and dental appearance provided substantial functional and psychological benefit.

Keywords: Corticobasal® implants; cortical anchorage; bimaxillary dental protrusion; pre-implant bone reduction; immediate functional loading; full-mouth rehabilitation; dento-facial esthetics; psychosocial impact of malocclusion.

1. Introduction

Bimaxillary dental protrusion and anterior positioning of the maxillary and mandibular dentoalveolar segments beyond the ideal skeletal profile is a common cause

of lip incompetence, an unesthetic convex facial profile, and significant psychosocial burden. Patients with this condition frequently report being subjected to teasing, mockery, or bullying regarding their appearance, which can profoundly affect self-esteem, social confidence, and quality of life. When bimaxillary protrusion is compounded by loss of posterior teeth, patients face a combined esthetic and functional deficit that is considerably more difficult to manage.

The conventional approach to correcting dentoalveolar protrusion relies on comprehensive orthodontic treatment, frequently combined with premolar extractions or, in more severe skeletal cases, orthognathic surgery, followed sequentially by implant placement to restore missing posterior teeth once orthodontic movement is complete and healing has occurred. This staged approach can require several years to complete and imposes a substantial cumulative burden, which may be poorly tolerated by patients experiencing acute psychological distress related to their appearance. It should be mentioned here also that earlier treatments with fixed orthodontic appliances have found to be the reason for

generalized and early tooth loss as well as indication for complete implants rehabilitation due to that orthodontic treatment^{10,11,12}. Considering the severe side effects, an orthodontic treatment would have been no option for this patient.

Cortical and basal implants offer a fundamentally different treatment paradigm. Rather than repositioning teeth orthodontically, the protrusive alveolar bone segment itself can be surgically reduced at the time of tooth extraction, correcting the dentoalveolar profile directly, while immediately loadable basal implants are anchored into the underlying dense cortical bone, a region that is resistant to resorption and is not compromised by the protrusive or resorbed crestal bone above it. This combined surgical-prosthetic approach allows correction of both the esthetic deformity and the missing dentition within a single, markedly compressed treatment episode.

This case report documents the full-mouth rehabilitation of a 57-year-old female patient with severe bimaxillary dental protrusion, bilateral posterior tooth loss in both arches, and substantial psychological distress attributed to bullying from family members and those around her.

Treatment combined extraction of the remaining compromised dentition, surgical resection of approximately 7 mm of protrusive alveolar bone in each jaw, and immediate placement eighteen basal implants in total, followed by immediate provisional loading and delivery of the final fixed prostheses.

2. Diagnosis: Clinical And Radiographic Assessment

2.1 Patient Profile

The patient was a 57-year-old female who presented with chief complaints of severe protrusion of the upper and lower anterior teeth, an unesthetic facial profile, difficulty chewing due to the loss of multiple posterior teeth in both jaws, and marked psychological distress. The patient reported that she had been repeatedly bullied and mocked by family members and by people around her because of her dental and facial appearance, and she expressed a strong wish for rapid, comprehensive correction. Her medical history was unremarkable, with no identified contraindications to extraction, alveolar surgery, or implant placement.

2.2 Extraoral and Intraoral Examination

Extraoral examination revealed marked bimaxillary dental protrusion with incompetent lip seal at rest and a convex facial profile.




Fig. 1: Pre-operative extraoral frontal photograph showing lip incompetence and dental / facial appearance secondary to bimaxillary dental protrusion.



Fig. 2: Pre-operative extraoral lateral (profile) photograph showing a convex facial profile and protrusive lip line.

Intraoral examination confirmed severely proclined and irregularly spaced maxillary and mandibular anterior teeth, with generalized diastemas, occlusal disharmony, and bilateral loss of posterior teeth in both the maxilla and the mandible. Remaining teeth demonstrated variable mobility, and at least one grossly carious, non-restorable molar was identified.



Fig. 3: Pre-operative intraoral photograph – frontal occlusal view showing bimaxillary anterior protrusion, irregular spacing, and posterior edentulous segments.



Fig. 4: Pre-operative intraoral photograph – anterior view showing midline diastema and proclined maxillary and mandibular incisors.



Fig. 5: Pre-operative intraoral photograph – lateral view showing a grossly carious mandibular molar and posterior tooth loss.



Fig. 6: Pre-operative intraoral photograph – tongue-retracted view confirming posterior mandibular tooth loss and irregular remaining dentition.

The clinical findings confirmed a complex full-mouth rehabilitation case characterized by severe bimaxillary dentoalveolar protrusion, bilateral posterior tooth loss in both arches, compromised residual dentition, and a substantial esthetic and psychosocial burden.

2.3 Radiographic Assessment

Panoramic radiography (OPG) was performed. Radiographic findings included:

- Generalized horizontal bone loss in the posterior segments of both arches, with edentulous ridges bilaterally.

- A markedly procumbent anterior alveolar bone contour in both the maxilla and the mandible, consistent with the clinically observed dental protrusion.
- Periapical and periodontal changes associated with several of the remaining anterior teeth.
- Adequate cortical bone thickness in the basal (cortical) regions of both jaws including the pterygo-maxillary buttress region, the posterior mandibular cortex, and the symphyseal region confirming suitability for basal implant placement once the protrusive alveolar segment was reduced.

Diagnosis: Bimaxillary dentoalveolar protrusion with bilateral posterior edentulism, compromised anterior dentition, and associated esthetic and psychosocial impairment.



Fig. 7: Pre-operative panoramic radiograph showing partial edentulism with missing teeth in the upper and lower arches, existing restorations on the upper right premolar and upper left molar, and a tilted / malpositioned upper left molar requiring evaluation prior to treatment.

3. Treatment Plan

3.1 Treatment Rationale

Given the severity of the dentoalveolar protrusion, the bilateral loss of posterior teeth in both arches, and the patient's urgent request for a rapid esthetic and functional solution in view of her psychological distress, it was decided to avoid prolonged conventional orthodontic treatment and instead pursue a combined surgical reduction and immediate full-mouth basal implant rehabilitation. The key objectives were:

- Extraction of all remaining non-restorable and malpositioned teeth in both arches.

- Surgical resection of approximately 7 mm of the protrusive alveolar bone segment in each jaw to correct the dentoalveolar profile and expose a level cortical bed. Immediate provisional loading within 24 to 72 hours with a fixed prosthesis.
- Placement of multiple basal implants in both arches, utilizing dense cortical bone and bypassing the compromised, protrusive alveolar bone.
- Immediate provisional loading with a fixed metal framework.
- Final full-arch prosthetic rehabilitation restoring an ideal occlusal relationship and a corrected facial profile.

No orthodontic tooth movement, sinus lifting, or staged bone grafting was planned.

3.2 Surgical Protocol

After completion of the required medical evaluation and confirmation that the patient had no contraindications to surgical intervention, the procedure was performed under local anesthesia. The remaining maxillary teeth were extracted, and approximately 7 mm of the protrusive maxillary alveolar bone was

then resected using a surgical bur / piezo device to correct the procumbent alveolar contour and to expose a level, dense cortical bed suitable for implant anchorage.




Fig. 8: Intra-operative photograph immediately following extraction and osteotomy, showing the resected alveolar ridge with active bleeding.




Fig. 9: Intra-operative photograph – close-up of the prepared alveolar ridge after bone resection, in preparation for basal implant placement.

Ten immediate basal implants were placed in the maxilla, engaging the dense cortical bone exposed by the alveolar resection. The identical surgical sequence extraction, resection of approximately 7 mm of protrusive alveolar bone, and implant placement was then carried out in the mandible, where six polished cortical and basal implants (BCS®, Manufacturer: Dr. Ihde Dental AG, Switzerland) were placed in addition to the two conical compression screws (TPG® uno, Manufacturer: Dr. Ihde Dental AG, Switzerland).

Eighteen cortical and basal implants in total were inserted across both arches. The extracted teeth and the resected segments of alveolar bone were retained for documentation.



Fig. 10: Photograph of the extracted teeth together with the resected segment of alveolar bone (approximately 7 mm), retrieved during ridge reduction.




Fig. 11: Close-up intra-operative photograph of the resected alveolar bone segment held with surgical forceps, illustrating the extent of ridge reduction.



Fig. 12: Post-operative panoramic radiograph of the mandible showing a full-arch implant-supported fixed prosthesis (bridge) with multiple dental implants placed bilaterally in the posterior and anterior regions, while the upper arch shows natural anterior teeth with no implants.



Fig. 13: Intra-operative photograph showing the immediate basal implant abutments in place with sutures, prior to framework try-in.

3.3 Metal Framework Try-In and Occlusal Adjustment

A working impression was recorded to capture the implant positions, and a master model incorporating implant analogs was fabricated to guide construction of the immediate metal framework.



Fig. 14: Working impression showing implant transfer components and analogs, used for fabrication of the immediate metal framework.

The metal framework was then seated over the implant abutments in both arches, and the accuracy of fit, passive insertion, and inter-arch relationship were carefully evaluated. Bite registration was taken for both the maxillary and mandibular arches to guide the final prosthetic arrangement, correction of the dental protrusion, and occlusal adjustment.

A total of eighteen immediate basal implants were placed, ten in the maxilla and eight in the mandible. Implant type and length were selected according to the density and thickness of the available cortical bone at each site following alveolar resection. The distribution is summarized below.




Fig. 15: Post-operative panoramic radiograph showing bilateral full-arch implant-supported fixed prostheses in both the maxilla and mandible, with multiple dental implants placed throughout both arches supporting complete edentulous rehabilitation.

4. Results and Follow-Up

4.1 Immediate and Short-Term Outcomes

The surgical phase was completed uneventfully without intra-operative complications. Post-operative discomfort was managed with standard analgesics and antibiotics. All eighteen basal implants demonstrated satisfactory primary stability at the time of placement, allowing progression to immediate prosthetic rehabilitation. The provisional metal frameworks were fitted within a short interval and immediately restored a corrected anterior tooth position, improved lip support, and functional occlusion.

The patient reported immediate satisfaction with the visible correction of her facial profile and dental appearance, and expressed considerable relief regarding the psychological distress she had previously experienced. Post-operative healing showed progressive mucosal healing around all implant abutments, with no signs of infection or peri-implant complications.

4.2 Final Prosthetic Delivery

Following framework verification and bite registration, the final prosthetic teeth were fitted and clinically evaluated.

Occlusal assessment confirmed a stable maxillo-mandibular relationship, a corrected anterior tooth position with resolution of the previous protrusion, and restoration of full posterior support in both arches. The patient was instructed to begin using the prosthesis for mastication and gradual functional adaptation, and definitive prosthetic cementation was subsequently completed.



Fig. 16: Final clinical photograph showing a close-up smile view of the upper anterior teeth, demonstrating a natural-looking, well-aligned fixed prosthesis with good gingival health and esthetic outcome.



Fig. 17: Final clinical intraoral photograph with cheek retractors showing a full-arch fixed prosthesis in both the upper and lower jaws, demonstrating good occlusion, symmetry, and healthy surrounding soft tissue.



Fig. 18: Final clinical photograph – lateral (profile) view of the patient smiling, showing natural lip support and a harmonious facial profile with the final anterior dental restoration in view.

4.3 Follow-Up

Follow-up examinations at one, three, and six months post definitive loading confirmed:

- Stable peri-implant soft tissue health with no signs of peri-implantitis.
 - Radiographically stable bone levels around all implant bodies.
 - No prosthetic complications (no fractures, no de-cementation, no screw loosening).
 - A markedly corrected dentofacial profile with resolution of the pre-treatment bimaxillary protrusion.
 - Patient-reported outcomes: high satisfaction with function, esthetics, and a substantial improvement in self-confidence and psychological wellbeing, with resolution of the distress previously associated with bullying.
- No sinus floor elevation procedure required.
 - No staged bone grafting or augmentation performed.
 - Immediate provisional loading of all eighteen implants.
 - Markedly shortened total treatment time compared with orthodontic-surgical implant sequencing.
 - Significant improvement in facial esthetics and psychological wellbeing.

4.4 Summary of Clinical Advantages

- Single-episode surgical correction of severe bimaxillary dental protrusion without conventional orthodontic treatment.
- Simultaneous rehabilitation of bilateral posterior edentulism in both arches.

5. Discussion

Bimaxillary dental protrusion combined with posterior tooth loss represents a particularly challenging clinical scenario, as it requires simultaneous correction of the dentoalveolar profile and reconstruction of missing posterior support. Conventional management and comprehensive orthodontic treatment, often combined with premolar extractions, followed sequentially by implant placement, can require an extended treatment period that may be poorly tolerated by patients experiencing significant psychosocial distress related to their appearance, as was reported in this case.

Surgical resection of the protrusive alveolar segment at the time of extraction allows direct correction of the dentoalveolar profile without relying on orthodontic tooth movement. When combined with immediately loadable basal implants anchored into the underlying dense cortical bone, this approach allows both the esthetic deformity and the missing dentition to be addressed within a single, markedly compressed treatment episode.

The decision to resect approximately 7 mm of alveolar bone in each jaw was based on the degree of protrusion observed clinically and radiographically, and was performed to expose a level, dense cortical bed capable of providing high primary implant stability. High primary stability achieved through cortical anchorage enabled immediate splinting of the implants with a fixed provisional restoration, consistent with the immediate loading protocols described for basal (cortical) implant systems.

From a patient-centered perspective, the psychosocial impact of visible dental protrusion, including the bullying reported by this patient from family members and those around her, should not be un-

derestimated. The ability to deliver a substantially corrected dentofacial appearance together with a fixed, functional dentition within a compressed treatment timeline offers considerable psychological benefit in addition to the functional and esthetic outcomes achieved.

The patient had developed a compensatory habitual smile pattern due to her previous severe dentoalveolar and skeletal protrusion. Before surgery, she habitually forced her upper lip to cover her prominent teeth and protrusive maxillo-mandibular profile. Despite successful orthognathic correction, this compensatory lip posture has persisted as a learned behavioral habit and remains evident during smiling.

6. Conclusion

This case report demonstrates that combining a full extraction in both jaws with vertical surgical resection of a protrusive alveolar segment with immediate Corticobasal® implant placement is a viable approach for the simultaneous correction of severe bimaxillary dental protrusion and rehabilitation of bilateral posterior edentulism in both arches.

The technique achieved a corrected dentofacial profile, restored full-arch function at an adequate vertical dimension, and provided substantial psychological benefit within a markedly shortened treatment timeline compared with conventional orthodontic-surgical implant sequencing. The six-month follow-up data confirm stable peri-implant tissues, prosthetic integrity, and high patient satisfaction.

Careful case diagnostics regarding the bi-alveolar protrusion, precise surgical execution of the alveolar reduction, and thorough prosthetic planning remain critical to the long-term success of this combined approach. Continued follow-up, typically after three months and then every twelve months (the so-called Belgrade protocol), will guarantee the predictability of the treatment modality shown here. Long-term observations give proof that this treatment and following the follow-up protocol strictly yields stable results with zero losses of BIPS® after year four. Corticobasal® implants undergo a dynamic interaction with the peri-implant bone and therefore, these implants will potentially stay in and functional forever⁸.

This new implant technology shows optimal long-term results, it avoids the hassles of peri-implantitis as well as bone augmentations and all kind of healing times. This technology is today the specialist standard, the patients' first method of choice, and it replaces the formerly used technology of "osseointegration"¹³.

Implant No.	Jaw	Region / Tooth Position	Implant Type	Length (mm)
1	Maxilla	Floor of the nose	BCS®	20*3
2	Maxilla	Floor of the nose	BCS®	20*3
3	Maxilla	Floor of the nose	BCS®	20*3
4	Maxilla	Floor of the nose	BCS®	20*3
5	Maxilla	Floor of the nose	BCS®	20*3
6	Maxilla	Floor of the nose	BCS®	20*3
7	Maxilla	Floor of the nose	BCS®	14*4.5
8	Maxilla	Floor of the nose	BCS®	14*4.5
9	Maxilla	Sphenoid bone	BCS®	20*4.5
10	Maxilla	Sphenoid bone	BCS®	20*4.5
11	Mandible	Chin	BCS®	20*3
12	Mandible	Chin	BCS®	20*3
13	Mandible	Chin	TPG® uno	19*3.5
14	Mandible	Chin	TPG® uno	15*3.5
15	Mandible	Mylohyoid region	BCS®	17*3.5
16	Mandible	Mylohyoid region	BCS®	17*4.5
17	Mandible	Mylohyoid region	BCS®	14*3.5
18	Mandible	Mylohyoid region	BCS®	14*4.5

Table 1: Distribution and characteristics of the immediate basal implants placed in the maxilla and mandible.

References

- 1** Ihde S, Sipic O. Immediate loading of basal implants: treatment protocol and overview of clinical results. *J Osseointegr.* 2009;1(2):49-54.
- 2** Scortecchi G, Misch CE, Benner KU. *Implants and Restorative Dentistry.* London: Martin Dunitz; 2001.
- 3** Peñarrocha-Diago M, Candel-Marti E, Ata-Ali J, Peñarrocha-Oltra D. Rehabilitation with tilted implants: a systematic review. *J Oral Implantol.* 2014;40(2):215-222.
- 4** Agliardi E, Panigatti S, Clericò M, Villa C, Malò P. Immediate rehabilitation of the edentulous jaws with full fixed prostheses supported by four implants: interim results of a single cohort prospective study. *Clin Oral Implants Res.* 2010;21(5):459-465.
- 5** Albrektsson T, Zarb G, Worthington P, Eriksson AR. The long-term efficacy of currently used dental implants: a review and proposed criteria of success. *Int J Oral Maxillofac Implants.* 1986;1(1):11-25.
- 6** Misch CE. *Dental Implant Prosthetics.* 2nd ed. St. Louis: Elsevier Mosby; 2015.
- 7** Proffit WR, Fields HW, Sarver DM. *Contemporary Orthodontics.* 5th ed. St. Louis: Elsevier Mosby; 2013.
- 8** Ihde S, Sipic O, Ihde A, Majeed, M M. Long-Term Outcomes of Immediately Loaded Cortical and Basal Implants in Full-Arch Rehabilitation: A Prospective Practice-Based Study, Experience-based and evidence-oriented Corticobasal® Implantology (EECI), Vol. 19, No. 2, International Implant Foundation Publishing, 2025
- 9** Seehra J, Fleming PS, Newton T, DiBiase AT. Bullying in orthodontic patients and its relationship to malocclusion, self-esteem and oral health-related quality of life. *J Orthod.* 2011;38(4):247-256.
- 10** Ihde S, Sipic O, Ihde A, Zappa U. Critical Appraisal of Traditional Dentistry – Part 3: Fixed Orthodontic Treatment: Functional and Biological Risks, Reasons for Failure and Hidden Long-Term Consequences, as Well as Ethical, Legal, and Social Considerations, Experience-based and evidence-oriented Corticobasal® Implantology (EECI), Vol. 19, No. 7, International Implant Foundation Publishing, 2025
- 11** Ihde A., Ihde S., Sipic O., Ihde A., Zappa U. Beschreibung des Ursprungs schwerer Schäden im Kausystem eines erwachsenen Patienten nach kieferorthopädischer Behandlung in der Jugend, sowie Beseitigung der Schäden durch Entfernung der gesamten Dentition und sofortige Rekonstruktion unter Verwendung der Technologie des Strategic Implant®, *Journal of Unwanted Results and Their Correction (JUR)*, Vol. 19, No. 2, International Implant Foundation Publishing, 2025 (article in German language)
- 12** Sipic O., Ihde A., Ihde S. A Case of Full-Mouth Rehabilitation Using Immediately Loaded Fixed Prostheses Supported by Corticobasal® Implants as Consequence of a Standardized Fixed Orthodontic Treatment, *Journal of Unwanted Results and their Correction (JUR)*, Vol. 18, No. 1, International Implant Foundation Publishing, 2024

13 Ihde S, Sipic O, Ihde A, Majeed, M M. A. Multi-Criteria Framework for Comparing Dental Implant Technologies: A 9-Field Decision Matrix Approach, Experience-based and evidence-oriented Corticobasal® Implantology (EECI), Vol. 19, No. 2, International Implant Foundation Publishing, 2025

ADVERTISEMENT

Previously

SIMPLADENT®

Three Days Later



Only within the scope of a complete restoration with Corticobasal® / BCS® implants was it possible to successfully correct all parameters of the masticatory function (Strategic Reset®).

SINGLE-PIECE IMPLANT

IMMEDIATE LOADING DENTAL SYSTEM

BCS®

CORTICOBASAL® IMPLANT

More information



www.implant.com



IHDEDENTAL 


ADVERTISEMENT

2015

SIMPLADENT®

MONTENEGRO

2025



This patient case was treated in 2015 at the Simpladent® clinic in Budva, Montenegro (www.simpladent.me). The bone level around the implants is completely unchanged after 10 years. KOS® compression screws and Corticobasal® BCS® implants were used. Vertical bone growth can be observed in both maxillary sinuses without the need for a sinus lift. This bone growth resulted from the increasing force applied to the jawbone via the implants.

SINGLE-PIECE IMPLANT

IMMEDIATE LOADING DENTAL SYSTEM

TPG® UNO

CORTICOBASAL® IMPLANT

More information



www.implant.com



IHDEDENTAL



YOU ARE GOING TO SEE THE LIGHT!

**100 % of our course
participants understand why
conventional dentistry as such
is a very doubtful occupation.**

We will show you how to stop the breakdown of the masticatory system by doing the „Strategic Reset®“ on modern implants. You will see and understand why natural dentitions break down, and why the results of the Strategic Reset® on modern Implant Corticobasal® implants are lasting.

ABOUT US



Since 2006, the International Implant Foundation (IF®) in Munich, Germany, has been at the forefront of implantology, advancing the field through innovative research and education. Our mission is to provide world-class implantology training for dentists, to support research and continuous exchange of knowledge & experience between professionals, and to inform the general public about the possibilities of modern oral implantology

16 Advantages of Osseofixation

Patients will hold you as their treatment provider, because you offer these advantages to them:

- | | | | |
|---|---|--|---|
| 1 Saves costs by 30-40% |  | 9 Aesthetic solutions for all patients |  |
| 2 Reduces treatment time by 98% |  | 10 Uninterrupted intra-bony perfusion |  |
| 3 Efficient workflow saves chair-time |  | 11 Easy long-term maintenance |  |
| 4 Immediate functional loading |  | 12 No peri-implantitis |  |
| 5 Low complication rate |  | 13 No patient selection |  |
| 6 Simple straight forward treatment |  | 14 Put more implants |  |
| 7 Immediate implant placement |  | 15 Start treatment immediately |  |
| 8 Preserves bone elasticity |  | 16 Cost-effective implants |  |

AIOW CURRICULUM



Our All-in-One-Week Curriculum is an intense program designed for dentists to master **tooth-free dentistry**.

This course provides a solid foundation for future learning and patient treatments. With hands-on training and immediate application of skills, you'll be ready to safely implement the latest implantology techniques. Enroll in our advanced dental implants course today.

Requirements

A valid dental degree is required to enroll in our Corticobasal® implantology training program.

Who Should Attend

- Dentists and oral/maxillofacial surgeons interested in immediate functional loading
 - Prosthetic specialists
-

Features

- Instructions from experienced implantologists
 - Learn how to work without bone augmentation
 - Avoid peri-implantitis simply by choosing the right implant
 - Immediate implant placement
 - How to solve cases at all stages of atrophy
-

Course Duration

- A full & intense 7-day training program for modern implantology and directly associated subjects.
 - Become a certified implantologist in just one week.
-

Conventional Implantology



1 Inspection Diagnostic procedures Treatment plan

2_a **Surgery 1**
Tooth removal

2_b **Surgery 2**
Bone augmentation/sinus-lifting
(necessary in up to 80% of the cases)

2_c **Surgery 3**
Implant placement
(adequate bone healing provided)

2_d **Surgery 4**
Placement of gingiva former

2_e Impression taking

3 Trying of the bridge frame
(5-10 days after impression taking)

4 Delivery of bridge (**4-24 months**
after implant placement)

Total

Treatment duration: 4 - 24 Months
Number of appointments: 7 - 12

Strategic Implant®



Inspection
Diagnostic procedures
Treatment plan

1

Removal of teeth, Implant
placement, Impression
& Bite taking

2

**Step 1 and 2 may be done in
the same (first) appointment.*

Trying of a sample bridge and aes-
thetic & functional corrections
(if required) **0 - 1 days** after
implant placement

3

Delivery of bridge (**1 - 3 days**
after implant placement)

4

Control of occlusion and
mastication

5

Total

Treatment duration: 2 - 4 Days
Number of appointments: 4 - 5

AIOW TEACHERS



Prof. Dr. Stefan Ihde

Surgical & Prosthetic Specialist and
1st Class IF[®] Teacher



Prof. Dr. Vitomir Konstantinović

Professor of Maxillofacial
Surgery and Implantology,
Director of the Clinic for Max-
illofacial Surgery, 1st Class IF[®]
Teacher and Member of the IF[®]
Board



Prof. Dr. Olga Sipić

Prosthetic Specialist - Implantologist
and 1st Class IF[®] Teacher



Prof. Dr. Antonina Ihde

Prosthetic Specialist, 1st Class IF[®] Teacher,
and Head of Dental Implant Faculty.



Learn from 7 world-renowned professors

in implantology, each bringing extensive experience and expertise to your training. Our faculty includes leading experts in dental implants.

WORLD-
RENOVED
PROFESSORS



DT Sanela Lazinica

Dentist technician and specialized IF® teacher for the work on the Strategic Implant®



Prof. Dr. Aleksandar Lazarov

Surgical and prothetical specialist: 1st Class IF® Teacher, Member of the IF® Board



Prof. Dr. Yan Vares

Craniomaxillofacial surgeon, Head of Department, Oral Implantology, 1st Class IF® Teacher



Dr. Fodor Romulus Calin

Surgical and prosthetic specialist, 1st Class IF® Teacher



SUCCESS STORIES

And Impressions from the course



DR. IONUTS UNGUREAN

I got to know that this way of implantology is pretty good, I got so excited about it, that I quit my job at a hospital. And I took over a clinic which works only with these implants. Since then I don't do anything else!



DR. MIGUEL

It changed my world, because with the Strategic Implant® everything is permitted, anything is possible, you improve oral health of the patient in only 24 hours, and the really important part for me you don't have peri-implantitis. Prof. Ihde forever!



DR. HABIB RITHA

I am very lucky to have learned this course. Thank you Dr Ihde for your generosity and your modesty. Thank you all to the team!

There is no realistic alternative to modern Corticobasal® implants and its technology.

Get in Touch: Register now to our course for tooth-free dentistry and long lasting results!

A world map with a light blue background. The landmasses are in a darker blue. A red circle with the letters 'SRB' in white is positioned over the Balkan region, specifically over Serbia.

SRB

LECTURE ROOM:

Golsvordijeva 37, 11000 Belgrade,
Republic of Serbia

PHONE & WEBSITE:

+ 381 69 394 3788 (Whatsapp)
all-in-one-week.implantfoundation.org
implantfoundation.org



SEEING IS BELIEVING!